

SPECIFICATION

N-ACETYLGLUCOSAMINE DERIVATIVE AND ITS APPLICATION

5 Technical Field

The invention relates to an N-acetylglucosamine derivative, and a hyaluronic acid production-promoting agent or a skin external preparation containing the derivative. The invention provides a skin external
10 preparation capable of retaining the vital and moist skin.

Background Art

Hyaluronic acid is a high molecular weight polysaccharide having a high water retention capability and
15 has drawn attention as a very important extracellular matrix component in the skin (Shingo SAKAI, Shintaro INOUE, Metabolism of Hyaluronic Acid and Wrinkle Formation, Fragrance Journal, Fragrance Journal Ltd., vol. 26, no. 4, 49-58, Apr. 15, 1998).

20 Further, hyaluronic acid has been known to have many functions such as maintaining cells, retaining smoothness and suppleness of the skin, rendering the skin resistant to external force such as mechanical force and preventing the infection with pathogenic bacteria (BIO INDUSTRY, CMC Co.,
25 Ltd, vol. 8 no. 5, 66(346)-68(346), May 1, 1991).

On the other hand, it is reported that the intensity of the hyaluronic acid staining observed in the intercellular part of the epidermis decreases with aging (Ludger J.M. Meyer and Robert Stern, Age-Dependent Changes
30 of Hyaluronan in Human Skin, The Journal of Investigative Dermatology, The society for Investigative Dermatology, Inc, Vol. 102, No. 4, pp. 385-389, Apr. 1994), and that hyaluronic acid is hardly detected in the part affected by solar elastosis caused by ultraviolet radiation (Takuo
35 TSUJI, Physiological Aging of Skin: Difference from Photo-aging, Clinical Dermatology, Igaku-Shoin Ltd., Special

Edition, vol. 51, no. 5、53-57, Apr. 15, 1997) and accordingly it is considered that the drying and the deterioration of the vital properties and the elasticity of the skin are caused and consequently wrinkles are increased.

5 For the improvement of such conditions, cosmetics containing hyaluronic acid has been applied to keep the moisture retention capability of the skin surface; however, hyaluronic acid, which is a high molecular weight molecule, is hard to penetrate into the skin, and thus, the
10 fundamental improvement cannot be expected. Accordingly, it is highly expected to develop a substance which can fundamentally improve the skin function by increasing the hyaluronic acid synthesizing capability of the cells by themselves.

15 Retinoic acid has been known as a hyaluronic acid production-promoting substance in the epidermis. Retinoic acid originally exists in the epidermis and is a substance involved in the proliferation and the differentiation of the epidermal cells. However, retinoic acid has the skin
20 irritation properties. From this point of view, it is desired to find a hyaluronic acid production-promoting substance with which such problems can be avoided.


On the other hand, it has been reported that N-acetylglucosamine, which is a saccharide constituting
25 hyaluronic acid, can promote hyaluronic acid production in the cultured epidermal cells about 1.5-times at a concentration of 5 mmol/L independently from the cell proliferation (Fine Chemical, CMC Co., Ltd., vol. 30, no. 22, 5-11, Dec. 15, 2001). However, a high concentration of
30 N-acetylglucosamine is required to exert the hyaluronic acid production-promoting effect and accordingly, it is desired to develop a material which has an sufficient effect even at lower concentrations in order to apply in the broader fields such as the field of cosmetics and
35 pharmaceuticals.

Disclosure of the Invention

Accordingly, the object of the invention is to provide a hyaluronic acid production-promoting agent and a skin external preparation which are expected to retain the vital and moist skin and improve wrinkles by promoting hyaluronic acid production and have a higher effect than N-acetylglucosamine.

In view of the above state of the art, the present inventors have diligently studied seeking means of solving the conventional problems. Consequently, they have found that the particular compounds enumerated below can very easily and strongly promote the hyaluronic acid production in the epidermis and dermis, and thus has accomplished the present invention.

That is, the invention provides an N-acetylglucosamine derivative represented by the following formula (1), a skin external preparation containing the N-acetylglucosamine derivative, and a hyaluronic acid production-promoting agent containing the N-acetylglucosamine derivative as active ingredient:
formula (1)




wherein R^1 is a hydrogen atom or an alkyl group having 2 to 18 carbon atoms; R^2 , R^3 , and R^4 are hydrogen atoms or acyl groups having 2 to 18 carbon atoms and may be all the same or different from others; the steric structure at position 1 may be α or β ; provided that R^1 , R^2 , R^3 , and R^4 must not be all hydrogen atoms.

The invention provides preferably an N-acetylglucosamine derivative represented by the following formula (2) or (3), a skin external preparation containing the N-acetylglucosamine derivative, and a hyaluronic acid production-promoting agent containing the N-

acetylglucosamine derivative as active ingredient:


formula (2)



wherein R⁵ is an alkyl group having 2 to 18 carbon atoms;

- 5 R⁶ is a hydrogen atom or an acetyl group; and the steric structure at position 1 may be α or β : and

formula (3)




- wherein R⁷ is a hydrogen atom or an alkyl group having 2 to
10 18 carbon atoms; R⁸ is an acyl group having 2 to 18 carbon
atoms; and the steric structure at position 1 may be α or β .

Particular examples of the N-acetylglucosamine
derivative represented by the above-mentioned formula (1),
or (2) or (3) are those represented by the following


- 15 formulas (4) to (15):

formula (4)



wherein the steric structure at position 1 may be α or β ;


formula (5)




20

wherein the steric structure at position 1 may be α or β ;


formula (6)




wherein the steric structure at position 1 may be α or β ;
formula (7)




5 wherein the steric structure at position 1 may be α or β ;
formula (8)




wherein the steric structure at position 1 may be α or β ;
formula (9)




10 wherein the steric structure at position 1 may be α or β ;
formula (10)




15 wherein the steric structure at position 1 may be α or β ;
formula (11)




wherein the steric structure at position 1 may be α or β ;
formula (12)




5 wherein the steric structure at position 1 may be α or β ;
formula (13)



wherein the steric structure at position 1 may be α or β ;
formula (14)




10 wherein the steric structure at position 1 may be α or β ;
and
formula (15)



15 wherein the steric structure at position 1 may be α or β .


Further, the invention relates to a hyaluronic acid production-promoting agent and a skin external preparation containing the N-acetylglucosamine derivative represented by the following formula (16) as active ingredient:

formula (16)



wherein R⁹ is an acyl group having 2 to 16 carbon atoms and the steric structure at position 1 may be α or β .

5 Particular examples of the N-acetylglucosamine derivative represented by the above-mentioned formula (16) are those represented by the following formula (17) ;
formula (17)



10 wherein the steric structure at position 1 may be α or β .

Brief Description of the Drawings


Fig. 1 is a diagram showing the result of a hyaluronic acid-production promoting test (Test Example 1)
15 in epidermal cells using compounds produced in Production Examples 1 to 3 and 6 to 8. Fig. 2 is a diagram showing the result of a hyaluronic acid-production promoting test (Test Example 1) in epidermal cells using compounds produced in Production Examples 4 and 5. Fig. 3 is a
20 diagram showing the result of a hyaluronic acid-production promoting test (Test Example 2) in epidermal cells using 2-acetamido-1,3,4,6-tetra-O-acetyl-2-deoxy- β -D-glucopyranoside [the compound represented by the formula (17)] produced in Production Examples 9 to 12. Fig. 4 is a
25 diagram showing the result of a hyaluronic acid-production promoting test (Test Example 3) in dermal cells using compounds produced in Production Examples 1 and 6.

Best Mode for Carrying Out the Invention

30 An N-acetylglucosamine derivative to be used in the


present invention is represented by the following formula (1) or formula (16):

formula (1)



5 wherein R^1 is a hydrogen atom or an alkyl group having 2 to 18 carbon atoms; R^2 , R^3 , and R^4 are hydrogen atoms or acyl groups having 2 to 18 carbon atoms and may be all the same or different from others; the steric structure at position 1 may be α or β ; and provided that R^1 , R^2 , R^3 , and R^4 must
10 not be all hydrogen atoms:


formula (16)



wherein R^9 denotes an acyl group having 2 to 16 carbon atoms and the steric structure at position 1 may be α or β .


15 In this case, R^1 is a hydrogen atom or a straight-chain or branched alkyl group having 2 to 18 carbon atoms, preferably 4 to 16 carbon atoms, and most preferably 8 to 12 carbon atoms and may be saturated or unsaturated. R^2 , R^3 , and R^4 are hydrogen atoms or straight-chain or branched
20 acyl groups having 2 to 18 carbon atoms and may be all the same or different from others and preferably all hydrogen atoms or acetyl groups. The steric structure at position 1 of the pyranose ring may be α or β . However, R^1 , R^2 , R^3 , and R^4 must not be all hydrogen atoms.

25 The N-acetylglucosamine derivative represented by the above-mentioned formula (1) is preferably those which can be represented by the following formula (2) or (3):
formula (2)



wherein R^5 is an alkyl group having 2 to 18 carbon atoms;
 R^6 is a hydrogen atom or an acetyl group; and the steric
 structure at position 1 may be α or β ; and

5 formula (3)



wherein R^7 is a hydrogen atom or an alkyl group having 2 to
 18 carbon atoms; R^8 is an acyl group having 2 to 18 carbon
 atoms; and the steric structure at position 1 may be α or β .


10 In this case, R^5 is a straight-chain or branched
 alkyl group having 2 to 18 carbon atoms, preferably 4 to 16
 carbon atoms, and most preferably 8 to 12 carbon atoms and
 may be saturated or unsaturated. R^6 is a hydrogen atom or
 an acetyl group, preferably all being hydrogen atoms or
 15 acetyl groups. R^7 is a hydrogen atom or a straight-chain
 or branched alkyl group having 2 to 18 carbon atoms,
 preferably 4 to 16 carbon atoms, and most preferably 8 to
 12 carbon atoms and may be saturated or unsaturated. R^8 is
 a straight-chain or branched acyl group having 2 to 18
 20 carbon atoms, preferably 6 to 16 carbon atoms, most
 preferably 8 to 12 carbon atoms and may be saturated or
 unsaturated. R^9 is a straight-chain or branched acyl group
 having 2 to 16 carbon atoms, preferably 2 to 8 carbon atoms,
 most preferably 2 to 4 carbon atoms.

25 Further, in formulas (1) to (17), the steric
 structure at position 1 of the pyranose ring illustrated
 with the wavy line part may be α or β and their mixture may
 be used without causing any problem.


Particular examples of the N-acetylglucosamine
 30 derivative represented by the above-mentioned general

formula (2) or (3) may be those represented by the following formulae (4) to (15):


formula (4)



5 wherein the steric structure at position 1 may be α or β ;
formula (5)




wherein the steric structure at position 1 may be α or β ;
formula (6)




10


wherein the steric structure at position 1 may be α or β ;
formula (7)




wherein the steric structure at position 1 may be α or β ;
15 formula (8)




wherein the steric structure at position 1 may be α or β ;
formula (9)



wherein the steric structure at position 1 may be α or β ;
formula (10)



5 wherein the steric structure at position 1 may be α or β ;
formula (11)




wherein the steric structure at position 1 may be α or β .


These compounds can be synthesized using a known
10 glycosylation reaction. The outline of an exemplary
synthesis method is as follows. In the case of a compound
represented by formula (2) wherein R^6 is a hydrogen atom, a
mixture of α - and β -glycosides can be produced by
glycosylation of an alcohol with N-acetylglucosamine in the
15 presence of an acid catalyst, and the α and β can be
separated by using a silica gel column. Further, β -
glycoside can be produced as a single compound by using
oxazoline synthesis method. The compound represented by
formula (3) can be produced by heating and dissolving N-
20 acetylglucosamine or the compound represented by formula (2)
in a solvent, adding a variety of fatty acid halides or
anhydrides, and optionally a catalyst, and then carrying
out the reaction.

The compounds represented by formulas (12) to (15)
25 will be explained below:

Formula (12)




wherein the steric structure at position 1 may be α or β ;
formula (13)




5

wherein the steric structure at position 1 may be α or β ;
formula (14)



10

wherein the steric structure at position 1 may be α or β ;
and
formula (15)




wherein the steric structure at position 1 may be α or β .

These compounds are commercially available 2-
15 acetamide-1,3,4,6-tetra-O-acetyl-2-deoxy- β -D-
glucopyranoside [the compound represented by the formula
(17)] or can be produced from this compound by
derivatization using glycosylation reaction according to
known oxazoline synthesis methods.

20

Particular examples of the N-acetylglucosamine
derivative represented by the formula (16) include those
which can be represented by the following formula (17):
formula (17)



wherein the steric structure at position 1 may be α or β .

The formulation amount of the N-acetylglucosamine derivative in a hyaluronic acid production-promoting agent and a skin external preparation is preferably 0.00001 to 5.0% by mass, more preferably 0.001 to 3.0% by mass, and most preferably 0.01 to 1.0% by mass on the basis of the total weight of the composition. If it is within the range, the aimed effect of the invention can sufficiently be exhibited.

The hyaluronic acid production-promoting agent and the skin external preparation of the present invention may be in a variety of application forms such as an ointment, lotion, emulsion, milk, cataplasm, pack, mist, foam, granule, powder, gel or the like. In the present invention, the skin external preparation is a preparation to be applied for external use to any skin of the body including the scalp and includes a bath agent. The base is not particularly limited as long as it is a base to be commonly used for external application. The final form of the agent may be cosmetics, pharmaceuticals, and quasi drugs.

The hyaluronic acid production-promoting agent and the skin external preparation of the present invention may suitably contain the following additives without departing from the scope of attaining the object of the present invention, in addition to those described above: tar type color additives; silicone oils such as dimethylpolysiloxane, methylphenylpolysiloxane, and cyclic silicone; carotenoid type coloring elements such as lutein, astaxanthin and fucoxanthin; coloring pigments such as iron oxide; preservatives such as parabens and phenoxyethanol; hydrocarbons such as paraffin and vaseline; vegetable oils such as olive squalane, rice squalane, rice germ glycerides,

jojoba oil, castor oil, safflower oil, olive oil, macadamia nut oil, and sunflower oil; waxes such as beeswax, Japan wax (*Rhus succedanea* fruit wax), and carnauba wax; ester oils such as octyldodecyl myristate, cetyl palmitate, 5 isostearyl isostearate, and isopropyl myristate; lower alcohols such as ethanol; higher alcohols such as cetanol, behenyl alcohol, stearyl alcohol, and a branched long chain aliphatic alcohol; sterols and their derivatives such as cholesterol, phytosterol, branched fatty acid cholesterol 10 ester, and macadamia nut fatty acid phytosteryl ester; processed oils such as hydrogenated oil; higher fatty acids such as stearic acid, myristic acid, isostearic acid, oleic acid, iso-type long chain fatty acid, and anteiso long chain fatty acid; terpenes such as limonene and 15 hydrogenated bisabolol; triglycerides such as glyceryl tricaprylcaprate, glyceryl 2-ethylhexanoate, glyceryl triiso-type long chain fatty acid ester, and glyceryl tripalmitate; anionic surfactants such as sodium cetylsulfate and N-stearoyl-L-glutamic acid salt; nonionic 20 surfactants such as polyoxyethylene alkyl ether, polyoxyethylene fatty acid ester, polyoxyethylene polyhydric alcohol fatty acid ester, polyoxyethylene hydrogenated castor oil, polyhydric alcohol fatty acid ester, modified silicone such as polyoxyethylene-modified 25 silicone, polyglycerin fatty acid ester, and sucrose ester; cationic surfactants such as tetraalkylammonium salt; amphoteric surfactants such as betaine type, sulfobetaine type, and sulfoamino-acid surfactants; natural type surfactants such as lecithin, lysophosphatidylcholine, 30 ceramide, and cerebroside; pigments such as titanium oxide and zinc oxide; antioxidants such as dibutylhydroxytoluene; inorganic salts such as sodium chloride, magnesium chloride, sodium sulfate, potassium nitrate, sodium sulfate, sodium metasilicate, and calcium chloride; organic acids and their 35 salts such as sodium citrate, potassium acetate, sodium succinate, sodium asparaginate, sodium lactate,

dichloroacetic acid, mevalonic acid, and glycyrrhizinic acid; organic amines and their salts such as ethanol amine hydrochloride, ammonium nitrate, arginine hydrochloride, diisopropylamine salt, urea, and decarboxycarnosine;

5 chelating agents such as edetic acid; thickeners such as xanthane gum, carboxyvinyl polymer, carrageenan, pectin, alkyl-modified carboxyvinyl polymer, and agar; neutralizing agents such as potassium hydroxide, diisopropanolamine, and triethanolamine; ultraviolet absorbents such as

10 hydroxymethoxybenzophenonesulfonate salt; polyhydric alcohols such as dipropylene glycol, marvitol, 1,3-butylene glycol, glycerin, propylene glycol, sorbitol, diglycerin, and raffinose; vitamins such as various amino acids, ascorbic acid, biotin, and tocopherol; and vitamin

15 derivatives such as ascorbic acid sulfate ester salt, ascorbic acid phosphate ester salt, and tocopherol nicotinate.

Further, by suitably formulating the following additives to the extent without departing from the scope of

20 attaining the object of the invention, the long-lasting effects of keeping the skin vital and moist and the higher anti-wrinkle effects are obtained: dermal hyaluronic acid production-promoting agents such as N-methyl-L-serine and yeast extract; hyaluronic acid decomposition suppressing

25 agents such as Naematoloma subliteritium extract, Boletopsis leucomelas extract, Mokkin (*Hibiscus syriacus*) extract, *Uncaria gambir* extract, and *Eugenia caryophyllus* flower extract; differentiation promoters of keratinocytes such as diisopropylamine dichloroacetate, niacin, mevalonic acid,

30 hot spring water, sodium metasilicate, and homogenized fruit; and barrier strengthening agents such as β -hydroxy- γ -aminobutyric acid and mevalonic acid.

Examples

35 The present invention will be illustrated below in more detail by the following examples. However, the

invention is not by anyway to be limited to the following examples.

(1) Production Examples of N-acetylglucosamine derivative

5 Production Example 1

Production of octyl(2-acetamido-2-deoxy) β -D-glucopyranoside [the compound represented by the formula (4)]:

After 2 g of 2-acetamido-1,3,4,6-tetra-O-acetyl-2-deoxy- β -D-glucopyranoside was dissolved in 20 mL of
10 anhydrous chloroform, 1.0 mL of trimethylsilyl trifluoromethanesulfonate was added and the mixture was stirred at room temperature for 5 hours. Chloroform was added to the reaction mixture, and after washing with an aqueous saturated sodium hydrogen carbonate solution, the
15 chloroform layer was dried over anhydrous magnesium sulfate and the solvent was distilled off under reduced pressure. The residue was dissolved in 15 mL of dichloroethane, and 0.89 mL of 1-octanol and 119 mg of (\pm)-camphor-10-sulfonic acid were added. The mixture was stirred at 60°C for 2
20 hours. Chloroform was added to the reaction mixture, and after washing with an aqueous saturated sodium hydrogen carbonate solution, the chloroform layer was dried over anhydrous magnesium sulfate, and the solvent was distilled off under reduced pressure. Finally, the resulting residue
25 was subjected to silica gel column chromatography (elution solvent; n-hexane : ethyl acetate = 2 : 3) to isolate a purified material. The obtained material was dissolved in a mixed solvent of 10 mL of methanol and 5 mL of 1,4-dioxane and a catalytic amount of 28% sodium methylate-methanol solution was added, and the mixture was stirred at
30 room temperature for 1 hour. After the reaction mixture was neutralized, the solvent was distilled off. Finally, the resulting residue was crystallized from water to obtain 840 mg of octyl(2-acetamido-2-deoxy) β -D-glucopyranoside as
35 a white crystal.

¹H-NMR measurement results of octyl(2-acetamido-2-

deoxy) β -D-glucopyranoside are as follows:

NMR (DMSO- d_6) δ : 0.85 (t, 3H, $J=6.6$ Hz), 1.23 (s, 10H), 1.40-
1.45 (m, 2H), 1.77 (s, 3H), 3.00-3.10 (m, 2H), 3.20-3.50 (m, 4H), 3.65-
5 3.75 (m, 2H), 4.25 (d, 1H, $J=8.3$ Hz), 4.40 (t, 1H), 4.78 (d, 1H), 4.87 (d,
1H), 7.58 (d, 1H, $J=8.7$ Hz).

Production Example 2

Production of 2-acetamido-2-deoxy-6-O-octanoyl- α -D-
10 glucopyranose [the compound represented by the formula
(5)]:

To 0.5 g of N-acetylglucosamine, 5 mL of pyridine
and 5 mL of N,N-dimethylformamide were added, and the
mixture was heated to 70°C under stirring, and 0.46 mL of
15 n-octanoyl chloride was dropwise added, and the reaction
was carried out for 4 hours. On completion of the reaction,
the reaction was extracted with ethyl acetate and washed
with 2 mol/L hydrochloric acid, and the obtained ethyl
acetate layer was dried over anhydrous magnesium sulfate
20 and then the solvent was distilled off under reduced
pressure. The resulting residue was purified by silica gel
column chromatography (elution solvent; chloroform:
methanol = 15 : 1) to obtain 170 mg of 2-acetamido-2-deoxy-
6-O-octanoyl- α -D-glucopyranose as a white crystal.

25 1 H-NMR measurement results of 2-acetamido-2-deoxy-6-
O-octanoyl- α -D-glucopyranose are as follows:

NMR (DMSO- d_6) δ : 0.92 (t, 3H, $J=6.8$ Hz), 1.33 (s, 10H), 1.55-
1.60 (m, 2H), 1.89 (s, 3H), 2.34 (t, 2H), 3.15-3.20 (m, 1H),
30 3.55-3.60 (m, 1H), 3.65-3.70 (m, 1H), 3.85-3.90 (m, 1H), 4.08
(dd, 1H, $J=6.0, 11.6$ Hz), 4.35 (dd, 1H, $J=2.1, 11.8$ Hz), 4.70
(d, 1H, $J=5.4$ Hz), 4.96 (t, 1H, $J=3.5, 4.3$ Hz), 5.13
(d, 1H, $J=5.8$ Hz), 6.54 (d, 1H, $J=4.7$ H), 7.61 (d, 1H, $J=8.1$ Hz).

35 Production Example 3

Production of octyl(2-acetamido-2-deoxy-6-O-octanoyl) β -D-

glucopyranoside [the compound represented by the formula (6)]:

In 1 mL of pyridine, 100 mg of the compound [the formula (4)] obtained in Production Example 1 was dissolved, and 61 μ L of n-octanoyl chloride was dropwise added, and the reaction was carried out for 4 hours. On completion of the reaction, after extracting with chloroform and washing with 2 mol/L hydrochloric acid, the ethyl acetate layer was dried over anhydrous magnesium sulfate and then the solvent was distilled off under reduced pressure. The resulting residue was purified by silica gel column chromatography (elution solvent; chloroform: methanol = 20 : 1) to obtain 40 mg of octyl(2-acetamido-2-deoxy-6-O-octanoyl) β -D-glucopyranoside as a white crystal.

1 H-NMR measurement results of octyl(2-acetamido-2-deoxy-6-O-octanoyl) β -D-glucopyranoside are as follows:

NMR (DMSO- d_6) δ : 0.85 (t, 3H, J=6.8Hz), 1.23(s, 18H), 1.40-1.45 (m, 2H), 1.50-1.55 (m, 2H), 1.78 (s, 3H), 2.28 (t, 2H), 3.05-3.10 (m, 1H), 3.25-3.40 (m, 4H), 3.60-3.65 (m, 1H), 4.05 (dd, 1H, J=7.2, 11.6Hz), 4.28 (d, 1H, J=8.0Hz), 4.30 (dd, 1H, J=1.6, 11.6Hz), 4.90 (d, 1H, J=4.8Hz), 5.12 (d, 1H, J=5.2Hz), 7.61 (d, 1H, J=8.4Hz).

25 Production Example 4

Production of butyl(2-acetamido-2-deoxy) β -D-glucopyranoside [the compound represented by the formula (7)]:

The same reaction was carried out as that of the Production example 1, except that 1-butanol was used in place of 1-octanol to obtain 280 mg of butyl(2-acetamido-2-deoxy) β -D-glucopyranoside as a white crystal.

1 H-NMR measurement results of butyl(2-acetamido-2-deoxy) β -D-glucopyranoside are as follows:

35 NMR (DMSO- d_6) δ : 0.83 (t, 3H, J=7.1Hz), 1.20-1.30 (m, 2H), 1.40-1.50 (m, 2H), 1.78(s, 3H), 3.00-3.05 (m, 2H), 3.25-3.45

(m, 4H), 3.65-3.70 (m, 2H), 4.26 (d, 1H, J=8.0Hz), 4.39 (t, 1H, J=5.8Hz), 4.77 (d, 1H, J=5.0Hz), 4.86 (d, 1H, J=4.4Hz), 7.57 (d, 1H, J=8.7Hz).

5 Production Example 5

Production of pentyl(2-acetamido-2-deoxy) β -D-glucopyranoside [the compound represented by the formula (8)]:

10 The same reaction was carried out as that of the Production Example 1, except that 1-pentanol was used in place of 1-octanol to obtain 150 mg of pentyl(2-acetamido-2-deoxy) β -D-glucopyranoside as a white crystal.

^1H -NMR measurement results of pentyl(2-acetamido-2-deoxy) β -D-glucopyranoside are as follows:

15

NMR (DMSO- d_6) δ : 0.85 (t, 3H, J=6.0Hz), 1.20-1.25 (m, 4H), 1.40-1.45 (m, 2H), 1.78 (s, 3H), 3.05-3.10 (m, 2H), 3.20-3.45 (m, 4H), 3.65-3.75 (m, 2H), 4.26 (d, 1H, J=8.0Hz), 4.40 (t, 1H, J=6.0Hz), 4.78 (d, 1H, J=4.8Hz), 4.87 (d, 1H), 7.58

20

(d, 1H, J=8.8Hz).

Production Example 6

Production of lauryl(2-acetamido-2-deoxy) β -D-glucopyranoside [the compound represented by formula (9)]:

25

The same reaction was carried out as that of the Production example 1, except that 1-dodecanol was used in place of 1-octanol to obtain 450 mg of lauryl(2-acetamido-2-deoxy) β -D-glucopyranoside as a white crystal.

^1H -NMR measurement results of lauryl(2-acetamido-2-deoxy) β -D-glucopyranoside are as follows:

30

NMR (DMSO- d_6) δ : 0.85 (t, 3H, J=6.0Hz), 1.23 (s, 18H), 1.40-1.45 (m, 2H), 1.84 (s, 3H), 3.00-3.10 (m, 2H), 3.20-3.50 (m, 4H), 3.65-3.75 (m, 2H), 4.25 (d, 1H, J=8.0Hz), 4.40 (t, 1H, J=5.6Hz), 4.79 (d, 1H, J=5.2Hz), 4.85 (d, 1H, J=4.4Hz), 7.08 (d, 1H, J=8.8Hz).

35

Production Example 7

Production of 2-acetamido-2-deoxy-6-O-palmitoyl- α -D-glucopyranose [the compound represented by formula (10)]:

5 To 1 g of N-acetylglucosamine, 5 mL of pyridine and 15 mL of N,N-dimethylformamide were added, and the mixture was heated to 70°C under stirring and 1.37 mL of palmitoyl chloride was dropwise added, and the reaction was carried out for 4 hours. On completion of the reaction, after
10 extracting with ethyl acetate and washing with 2 mol/L hydrochloric acid, the ethyl acetate layer was dried over anhydrous magnesium sulfate and then the solvent was distilled off under reduced pressure. The resulting residue was purified by silica gel column chromatography
15 (elution solvent; chloroform: methanol = 15 : 1) to obtain 710 mg of 2-acetamido-2-deoxy-6-O-palmitoyl- α -D-glucopyranose as a white crystal.

^1H -NMR measurement results of 2-acetamido-2-deoxy-6-O-palmitoyl- α -D-glucopyranose are as follows:

20

NMR (DMSO- d_6) δ : 0.85 (t, 3H, J=6.5Hz), 1.25 (s, 24H), 1.45-1.55 (m, 2H), 1.82 (s, 3H), 2.30 (t, 2H), 3.05-3.15 (m, 1H), 3.45-3.65 (m, 2H), 3.75-3.85 (m, 1H), 4.00 (dd, 1H, J=5.7, 11.8Hz), 4.28 (dd, 1H, J=2.0, 11.8Hz), 4.65
25 (d, 1H, J=5.7Hz), 4.90 (t, 1H, J=3.7, 4.1Hz), 5.07 (d, 1H, J=5.7Hz), 6.45 (d, 1H, J=4.5H), 7.55 (d, 1H, J=8.1Hz).

Production Example 8

Production of geranyl(2-acetamido-2-deoxy) β -D-glucopyranoside [the compound represented by formula (11)]:
30

The same reaction was carried out as that of the Production Example 1, except that geraniol was used in place of 1-octanol to obtain 890 mg of geranyl(2-acetamido-2-deoxy) β -D-glucopyranoside as a white crystal.

35

^1H -NMR measurement results of geranyl(2-acetamido-2-deoxy) β -D-glucopyranoside are as follows:

NMR (DMSO- d_6) δ : 1.57, 1.60, 1.65 (3s, 9H), 1.79 (s, 3H), 1.95-
2.05 (m, 4H), 3.05-3.10 (m, 2H), 3.30-
3.40 (m, 2H), 3.47 (dt, 1H, $J=5.2, 12.0$ Hz), 3.68 (dd, 1H, $J=5.6, 11.6$ Hz
5), 4.02 (dd, 1H, $J=7.2, 12.0$ Hz), 4.17 (dd, 1H, $J=5.2$ Hz), 4.87 (d, 1H), 5.
05-5.10 (m, 1H), 5.21 (t, 1H, $J=6.4$ Hz), 7.59 (d, 1H, $J=8.8$ Hz).

Production Example 9

Production of ethyl(2-acetamido-3,4,6-tri-O-acetyl-2-
10 deoxy) β -D-glucopyranoside [the compound represented by
formula (12)]:

After 1 g of 2-acetamido-1,3,4,6-tetra-O-acetyl-2-
deoxy- β -D-glucopyranoside [the compound represented by
formula (17)] was dissolved in 10 mL of anhydrous
15 chloroform, 0.5 mL of trimethylsilyl
trifluoromethanesulfonate was added, and the mixture was
stirred at room temperature for 5 hours. Chloroform was
added to the reaction mixture, and after washing with an
aqueous saturated sodium hydrogen carbonate solution, the
20 chloroform layer was dried over anhydrous magnesium sulfate
and then the solvent was distilled off under reduced
pressure. The residue was dissolved in 8 mL of
dichloroethane, and 0.17 mL of ethanol and 60 mg of (\pm)-
camphor-10-sulfonic acid were added, and the mixture was
25 stirred at 60°C for 2 hours. Chloroform was added to the
reaction mixture, and after washing with an aqueous
saturated sodium hydrogen carbonate solution, the
chloroform layer was dried over anhydrous magnesium sulfate
and the solvent was distilled off under reduced pressure.
30 Finally, the resulting residue was crystallized by using
ether and n-hexane to obtain 0.5 g of ethyl(2-acetamido-
3,4,6-tri-O-acetyl-2-deoxy) β -D-glucopyranoside as a white
solid.

1 H-NMR measurement results of ethyl(2-acetamido-
35 3,4,6-tri-O-acetyl-2-deoxy) β -D-glucopyranoside are as
follows:

NMR (CDCl₃) δ : 1.18 (t, 3H, J=7.3Hz), 1.93, 2.00, 2.05
(4s, 12H), 3.56 (m, 1H), 3.68 (m, 1H), 3.78
(dt, 1H, J=8.4, 8.8, 10.4Hz), 3.88 (m, 1H), 4.12
5 (dd, 1H, J=2.4, 14Hz), 4.25 (dd, 1H, J=4.8, 12.4Hz), 4.69
(d, 1H, J=8.4Hz), 5.03 (t, 1H, J=9.6Hz), 5.30 (t, 1H, J=9.2Hz),
5.43 (d, 1H, J=8.8Hz).

Production Example 10

10 Production of pentyl(2-acetamido-3,4,6-tri-O-acetyl-2-deoxy) β -D-glucopyranoside [the compound represented by formula (13)]:

After 1.5 g of 2-acetamido-1,3,4,6-tetra-O-acetyl-2-deoxy- β -D-glucopyranoside was dissolved in 15 mL of
15 anhydrous chloroform, 0.75 mL of trimethylsilyl trifluoromethanesulfonate was added, and the mixture was stirred at room temperature for 5 hours. Chloroform was added to the reaction mixture, and after washing with an aqueous saturated sodium hydrogen carbonate solution, the
20 chloroform layer was dried over anhydrous magnesium sulfate and then the solvent was distilled off under reduced pressure. The residue was dissolved in 15 mL of dichloroethane and 0.51 mL of n-amyl alcohol and 99 mg of (\pm)-camphor-10-sulfonic acid were added thereto and the
25 mixture was stirred at 60°C for 2 hours. Chloroform was added to the reaction mixture, and after washing with an aqueous saturated sodium hydrogen carbonate solution, the chloroform layer was dried over anhydrous magnesium sulfate and the solvent was distilled off under reduced pressure.
30 Finally, the resulting residue was subjected to silica gel column chromatography (elution solvent; n-hexane : ethyl acetate = 2 : 3) to isolate and purify 1.1 g of pentyl(2-acetamido-3,4,6-tri-O-acetyl-2-deoxy) β -D-glucopyranoside as a white solid.

35 ¹H-NMR measurement results of pentyl(2-acetamido-3,4,6-tri-O-acetyl-2-deoxy) β -D-glucopyranoside are as

follows:

NMR (CDCl₃) δ : 0.87 (t, 3H, J=6.2Hz), 1.28 (s, 4H), 1.50-1.55 (m, 2H), 2.00, 2.03, 2.14 (4s, 12H), 3.45 (dt, 1H, J=7.2, 9.2Hz),
5 3.65-3.70 (m, 1H), 3.75-3.85 (m, 2H), 4.13 (dd, 1H, J=2.4, 12.4Hz), 4.25 (dd, 1H, J=4.8, 12.4Hz), 4.66 (d, 1H, J=8.0Hz), 5.03 (t, 1H, J=9.6Hz), 5.29 (t, 1H, J=9.2Hz), 5.40 (d, 1H, J=8.8Hz).

10 Production Example 11

Production of octyl(2-acetamido-3,4,6-tri-O-acetyl-2-deoxy) β -D-glucopyranoside [the compound represented by formula (14)]:

After 2 g of 2-acetamido-1,3,4,6-tetra-O-acetyl-2-deoxy- β -D-glucopyranoside was dissolved in 20 mL of
15 anhydrous chloroform, 1.0 mL of trimethylsilyl trifluoromethanesulfonate was added thereto and the mixture was stirred at room temperature for 5 hours. Chloroform was added to the reaction mixture, and after washing with
20 an aqueous saturated sodium hydrogen carbonate solution, the chloroform layer was dried over anhydrous magnesium sulfate and then the solvent was distilled off under reduced pressure. The residue was dissolved in 15 mL of dichloroethane, and 0.89 mL of n-octanol and 119 mg of
25 (\pm)-camphor-10-sulfonic acid were added, and the mixture was stirred at 60°C for 2 hours. Chloroform was added to the reaction mixture, and after washing with an aqueous saturated sodium hydrogen carbonate solution, the chloroform layer was dried over anhydrous magnesium sulfate
30 and the solvent was distilled off under reduced pressure. Finally, the resulting residue was subjected to silica gel column chromatography (elution solvent; n-hexane : ethyl acetate = 2 : 3) to isolate and purify to obtain 1.5 g of octyl(2-acetamido-3,4,6-tri-O-acetyl-2-deoxy) β -D-glucopyranoside as a white solid.
35

¹H-NMR measurement results of octyl(2-acetamido-

3,4,6-tri-O-acetyl-2-deoxy) β -D-glucopyranoside are as follows:

5 NMR (CDCl₃) δ : 0.86 (t, 3H, J=6.2Hz), 1.25 (s, 10H), 1.50-1.55 (m, 2H), 1.92, 1.99, 2.00, 2.05 (4s, 12H), 3.40-3.45 (m, 1H), 3.65-3.70 (m, 1H), 3.75-3.90 (m, 2H), 4.10 (d, 1H, J=12.2Hz), 4.23 (d, 1H, J=4.8, 12.2Hz), 4.65 (d, 1H, J=8.3Hz), 5.03 (t, 1H, J=9.8Hz), 5.28 (t, 1H, J=9.8Hz), 5.41 (d, 1H, J=8.7Hz).

10 Production Example 12

Production of geranyl(2-acetamido-3,4,6-tri-O-acetyl-2-deoxy) β -D-glucopyranoside [the compound represented by formula (15)]:

15 After 5 g of 2-acetamido-1,3,4,6-tetra-O-acetyl-2-deoxy- β -D-glucopyranoside was dissolved in 50 mL of anhydrous chloroform, 2.6 mL of trimethylsilyl trifluoromethanesulfonate was added, and the mixture was stirred at room temperature for 5 hours. Chloroform was added to the reaction mixture, and after washing with an
20 aqueous saturated sodium hydrogen carbonate solution, the chloroform layer was dried over anhydrous magnesium sulfate and the solvent was distilled off under reduced pressure. The residue was dissolved in 20 mL of dichloroethane, and 2.5 mL of geraniol and 298 mg of (\pm)-camphor-10-sulfonic
25 acid were added and the mixture was stirred at 60°C for 2 hours. Chloroform was added to the reaction mixture, and after washing with an aqueous saturated sodium hydrogen carbonate solution, the chloroform layer was dried over anhydrous magnesium sulfate and the solvent was distilled
30 off under reduced pressure. Finally, the resulting residue was subjected to silica gel column chromatography (elution solvent; n-hexane : ethyl acetate = 2 : 3) to isolate and purify to obtain 4.3 g of geranyl(2-acetamido-3,4,6-tri-O-acetyl-2-deoxy) β -D-glucopyranoside as a white solid.

35 ¹H-NMR measurement results of geranyl(2-acetamido-3,4,6-tri-O-acetyl-2-deoxy) β -D-glucopyranoside are as

follows:

NMR (CDCl₃) δ : 1.59, 1.64, 1.67 (3s, 9H), 1.91 (s, 3H), 1.99, 2.00, 2.05 (3s, 9H), 3.60-3.65 (m, 1H), 3.78
5 (dt, 1H, J=2.8, 12.4Hz), 4.10-4.30 (m, 3H), 4.68 (d, 1H, J=8.8Hz), 5.00-5.10 (m, 2H), 5.20-5.40 (m, 2H).

(2) Test Examples using N-acetylglucosamine derivatives

Next, Test Examples for evaluating the N-
10 acetylglucosamine derivatives produced in the above-mentioned Production Examples of the invention will be described.

Test Example 1 (Test for hyaluronic acid production
15 promotion in normal human epidermal cells)

Normal human epidermal cells (manufactured by Kurabo Industries Ltd.) were inoculated into 24-well plates and cultured in a culture medium for proliferation to the
20 confluent density. Then N-acetylglucosamine derivatives produced in the above-mentioned Production Examples 1 to 3 and 6 to 8 were added to a final concentration of 50 μ mol/L (25 μ mol/L, only for the derivative obtained in Production Example 7), and N-acetylglucosamine derivatives produced in
25 the Production Examples 4 and 5 were added to a final concentration of 1 mmol/L. After 48-hour culture from the addition, the hyaluronic acid released to the culture medium was measured. The measurement of the hyaluronic acid was carried out by using a commercially available
30 hyaluronic acid measurement kit (manufactured by Chugai Pharmaceutical Co., Ltd.).

The amount of hyaluronic acid produced by dermal fibroblast cells cultured in the culture medium containing a test substance is represented as a ratio relative to the
35 amount of hyaluronic acid produced by the cells cultured in the culture medium containing N-acetylglucosamine at a

final concentration of 1 mmol/L (which is defined as 1, Comparative Example 1). The results are shown in Fig. 1 and Fig. 2.

As shown in Fig. 1, octyl(2-acetamido-2-deoxy) β -D-glucopyranoside [the compound represented by formula (4)] of Production Example 1, 2-acetamido-2-deoxy-6-O-octanoyl- α -D-glucopyranose [the compound represented by formula (5)] of Production Example 2, octyl(2-acetamido-2-deoxy-6-O-octanoyl) β -D-glucopyranoside [the compound represented by formula (6)] of Production Example 3, lauryl(2-acetamido-2-deoxy) β -D-glucopyranoside [the compound represented by formula (9)] of Production Example 6, 2-acetamido-2-deoxy-6-O-palmitoyl- α -D-glucopyranose [the compound represented by formula (10)] of Production Example 7, and geranyl(2-acetamido-2-deoxy) β -D-glucopyranoside [the compound represented by formula (11)] of Production Example 8 exhibited 2.5- to 3.5- higher hyaluronic acid production-promoting effect at about 1/20 concentration (about 1/40 in the case of Production Example 7) when compared with N-acetylglucosamine. Especially, the compound represented by formula (3) wherein R⁷ is hydrogen was found having the high production-promoting activity.

Further, as shown in Fig. 2, butyl(2-acetamido-2-deoxy) β -D-glucopyranoside [the compound represented by formula (7)] of Production Example 4 and pentyl(2-acetamido-2-deoxy) β -D-glucopyranoside [the compound represented by formula (8)] of Production Example 5 exhibited 1.5- to 3- times higher hyaluronic acid production-promoting effect at the same concentration compared with N-acetylglucosamine.

Test Example 2 (Test for hyaluronic acid production promotion in normal human epidermal cells)

Normal human epidermal cells (manufactured by Kurabo Industries Ltd.) were inoculated into 24-well plates and

cultured in a culture medium for proliferation to the
confluent density, and then 2-acetamido-1,3,4,6-tetra-O-
acetyl-2-deoxy- β -D-glucopyranoside represented by formula
(17) to a final concentration of 100 μ mol/L, N-
5 acetylglucosamine derivative obtained in Production Example
9 at a final concentration of 1 mmol/L, and N-
acetylglucosamine derivatives obtained in Production
Examples 10 to 12 at a final concentration of 100 μ mol/L
were added. After 48-hour culture from the addition, the
10 amount of hyaluronic acid released into the culture medium
was measured. The measurement of the hyaluronic acid was
carried out by using a commercially available hyaluronic
acid measurement kit (manufactured by Chugai Pharmaceutical
Co., Ltd.). The results are shown in Fig. 3.

15 As shown in Fig. 3, ethyl(2-acetamido-3,4,6-tri-O-
acetyl-2-deoxy) β -D-glucopyranoside [the compound
represented by formula (12)] of Production Example 9
exhibited 6.5-times higher hyaluronic acid production-
promoting effect at the same concentration when compared
20 with N-acetylglucosamine. Further, pentyl(2-acetamido-
3,4,6-tri-O-acetyl-2-deoxy) β -D-glucopyranoside [the
compound represented by formula (13)] of Production Example
10, octyl(2-acetamido-3,4,6-tri-O-acetyl-2-deoxy) β -D-
glucopyranoside [the compound represented by formula (14)]
25 of Production Example 11, geranyl(2-acetamido-3,4,6-tri-O-
acetyl-2-deoxy) β -D-glucopyranoside [the compound
represented by formula (15)] of Production Example 12, and
2-acetamido-1,3,4,6-tetra-O-acetyl-2-deoxy- β -D-
glucopyranoside [the compound represented by formula (17)]
30 exhibited 3.8-, 2.8-, 3.7-, and 3.0-times higher hyaluronic
acid production-promoting effect at 1/10 concentration
compared with N-acetylglucosamine, respectively.

Test Example 3 (Test for hyaluronic acid production
35 promotion in normal human dermal fibroblast cells)

Normal human dermal fibroblasts (manufactured by American Type Culture Collection) were inoculated into 24-well plates and cultured in a culture medium for proliferation to be confluent and then the 100 $\mu\text{mol/L}$ of N-acetylglucosamine derivative in the above-mentioned
5 Production Example 1 and 25 $\mu\text{mol/L}$ of N-acetylglucosamine derivative in the Production Example 6 were added. After 48-hour culture from the addition, the amount of hyaluronic acid released into the culture medium was measured. The
10 measurement of the hyaluronic acid was carried out by using commercially available hyaluronic acid measurement kit (manufactured by Chugai Pharmaceutical Co., Ltd.).

The amount of hyaluronic acid produced by dermal fibroblast cells cultured in the culture medium containing
15 a test substance is represented as a ratio relative to the amount of hyaluronic acid produced by the cells in the culture medium containing N-acetylglucosamine at a final concentration of 1 mmol/L (Comparative Example 2). The results are shown in Fig. 4.

20 As shown in Fig. 4, both octyl(2-acetamido-2-deoxy) β -D-glucopyranoside [the compound represented by formula (4)] of Production Example 1 and lauryl(2-acetamido-2-deoxy) β -D-glucopyranoside [the compound represented by formula (9)] of Production Example 6
25 exhibited 1.5- to 2- times higher hyaluronic acid production-promoting effect at an approximately 1/10 to 1/40 concentration compared with N-acetylglucosamine.

Test Examples 4 to 7 and Comparative Examples 3 and 4
30 (Evaluation by subjects)

Women in forties to sixties as the subjects were divided into 8 groups each consisting of 20 women. Creams with the compositions shown in Table 1 (Test Examples 4 and 5 and Comparative Example 3) and face lotions with the
35 compositions shown in Table 2 (Test Examples 6 and 7 and Comparative Example 4) were each given to a separate group

and an appropriate amount was applied to the faces of the subjects twice a day for 3 consecutive months. After successive use, the vital feeling of the skin was evaluated.

The evaluation was carried out based on the following four grades: remarkably effective (the vital feeling of the skin was considerably improved), effective (the vital feeling of the skin was well improved), slightly effective (the vital feeling of the skin was improved), and no effect (no change). The effect was determined based on the ratio (%) of the total number of the subjects who evaluated the creams or the face lotions as being remarkably effective and effective.

Table 1

	Test Example 4	Test Example 5	Comparativ e Example 3
Compound represented by formula (4) (Production Example 1)	0.1	-	-
Compound represented by formula (17)	-	0.1	-
Stearic acid	2		
Monostearic acid glycerin	2		
Cetanol	3		
Cholesterol	0.5		
Vaseline	2		
Squalane	10		
Liquid paraffin	10		
Dimethylpolysiloxane	1		
Butylparaben	0.1		
Methylparaben	0.1		
Sodium N-stearoylglutamate	1		
Glycerin dipropylene glycol	5		
Purified water	balance		
Total	100		
Evaluation (%)	80	80	45

* The contents are all based on % by mass.

Table 2

	Test Example 6	Test Example 7	Comparativ e Example 4
Compound represented by formula (4) (Production Example 1)	0.01	-	-
Compound represented by formula (17)	-	0.01	-
Ethanol	10		
Polyoxyethylene hydrogenated castor oil (60 E.O.)	1		
Glycerin	3		
1,3-butylene glycol	2		
Dipropylene glycol	3		
Monopotassium phosphate	0.05		
Dipotassium phosphate	0.05		
Edetate disodium	0.05		
Methylparaben	0.1		
Purified water	balance		
Total	100		
Evaluation (%)	55	55	30

* The contents are all based on % by mass.

5 The results shown in Table 1 and Table 2 revealed that the cosmetics using N-acetylglucosamine derivatives of the invention had the effect of improving the vital feeling of the skin.

10. Additionally, in any case using the cosmetics of the Test Examples, no symptoms considered to be the side effects such as red rash, inflammation, and the like were observed on the skin and thus it was made clear that the cosmetics according to the invention were excellent also in the safety.

(3) Examples
15 Example 1 (Skin cream)

A skin cream with the following composition was prepared by a conventional method.

Ingredients	Formulation amount (% by mass)
Beeswax	2.0
Stearic acid	5.0
Stearyl alcohol	5.0
Hydrogenated lanolin	2.0
Squalene	20.0
Sorbitan monostearate	3.0
Polyoxyethylene (20) sorbitan monostearate	3.0
Propylene glycol	5.0
Methylparaben	0.2
Compound represented by formula (4) (Production Example 1)	0.1
Purified water	Balance in the total 100

Example 2 (Skin cream)

5 A skin cream with the following composition was
prepared by a conventional method.

Ingredients	Formulation amount (% by mass)
Beeswax	2.0
Stearic acid	5.0
Stearyl alcohol	5.0
Hydrogenated lanolin	2.0
Squalene	20.0
Sorbitan monostearate	3.0
Polyoxyethylene (20) sorbitan monostearate	3.0
Propylene glycol	5.0
Methylparaben	0.2
Compound represented by formula (5) (Production Example 2)	0.5
Purified water	Balance in the total 100

Example 3 (Skin lotion)

A skin lotion with the following composition was prepared by a conventional method.

Ingredients	Formulation amount (% by mass)
Olive oil	10.0
Isopropyl myristate	1.0
Polyoxyethylene (6) nonylphenyl ether	0.5
Propylene glycol	1.0
Glycerin	2.0
Methylparaben	0.1
Ethanol	7.0
Compound represented by formula (6) (Production Example 3)	0.5
Purified water	Balance in the total 100

5 Example 4 (Skin lotion)

A skin lotion with the following composition was prepared by a conventional method.

Ingredients	Formulation amount (% by mass)
Olive oil	10.0
Isopropyl myristate	1.0
Polyoxyethylene (6) nonylphenyl ether	0.5
Propylene glycol	1.0
Glycerin	2.0
Methylparaben	0.1
Ethanol	7.0
Compound represented by formula (8) (Production Example 5)	1.0
Purified water	Balance in the total 100

Example 5 (Bath agent)

10 A bathing agent with the following composition was prepared by a conventional method.

Ingredients	Formulation amount (% by mass)
Compound represented by formula (5) (Production Example 2)	1.0
Sodium hydrogen carbonate	balance
Sodium carbonate	20.0
Sodium sulfate	15.0
Sodium chloride	7.5
Silicic acid anhydride	0.5
1,3-butylene glycol	1.0
Urea	1.0
Seaweed extract	1.0
Color additive	q.s.
Dextrin	q.s.
Fragrance	q.s.

Examples 6 to 8 (Skin cream)

Skin creams with the following compositions were prepared by a conventional method.

Ingredients	Formulation amount (% by mass)		
	Example 6	Example 7	Example 8
Compound represented by formula (4) (Production Example 1)	0.1	-	-
Compound represented by formula (5) (Production Example 2)	-	0.1	-
Compound represented by general formula (6) (Production Example 3)	-	-	0.1
Stearic acid	1	1	-
Isostearic acid	-	-	1
Monostearic acid glycerin	2	2	2
Behenyl alcohol	2	2	2
Bleached beeswax	1	1	-
Cetyl myristate	1	1	1
Sorbitan sesquioleate	1	1	1

N-stearoylphytosphingosine	0.1	0.1	0.1
Hydrogenated lecithin	0.1	0.1	0.1
Plant squalane	5	5	5
Octyldodecyl myristate	5	5	5
Phellodendron bark extract	0.1	1	0.1
Pyracantha fortuneana fruit extract	0.1	0.3	-
Water-soluble glycyrrhiza extract	-	-	0.1
1,3-butylene glycol	5	10	5
Concentrated glycerin	5	5	5
p-oxybenzoic acid ester	0.2	0.2	0.2
N-acetylglucosamine oligomer	0.1	0.1	0.1
Ascorbic acid phosphate ester magnesium salt	0.1	0.1	0.1
Ascorbic acid phosphate ester sodium salt	0.1	0.1	0.1
γ-aminobutyric acid	0.1	0.1	0.1
Sodium N-stearoylglutamate	0.2	0.2	0.2
Alkyl-modified carboxyvinyl polymer *1	0.05	0.05	0.05
Nicotinic acid amide	0.1	0.1	0.1
Sarcosine	0.1	0.1	0.1
Purified water	balance	balance	balance

*1: PEMULEN TR-1, manufactured by B. F. Goodrich

Examples 9 to 11 (Lotion)

5 Lotions with the following compositions were prepared by a conventional method.

Ingredients	Formulation amount (% by mass)		
	Example 9	Example 10	Example 11
Compound represented by formula (4) (Production Example 1)	0.1	-	-
Compound represented by formula (5) (Production Example 2)	-	0.1	-
Compound represented by formula (6)	-	-	0.1

(Production Example 3)

Phellodendron bark extract	0.1	0.3	0.3
Hibiscus extract	0.2	0.5	0.5
Lactic acid bacteria culture medium	0.1	0.1	0.1
1,3-butylene glycol	5	5	5
Dipropylene glycol	5	5	5
Raffinose	1	1	1
Ethanol	1	1	1
Phenoxyethanol	0.2	0.2	0.2
Pectin	0.05	0.05	0.05
Xanthane gum	0.1	0.1	0.1
Sodium citrate	0.05	0.05	0.05
Equisetum arvense extract	0.1	0.1	0.1
Diisopropylamine dichloroacetic acid	0.2	0.2	0.2
γ -amino- β -hydroxybutyric acid	0.2	0.2	0.2
Sodium hyaluronate	0.001	0.001	0.001
Dipotassium glycyrrhizate	0.2	0.2	0.2
Naematoloma sublateritium extract	0.05	0.05	0.05
Decarboxycarnosine hydrochloride	0.05	0.05	0.05
Fragrance	0.02	0.02	0.02
Purified water	balance	balance	balance

Examples 12 to 14 (Gel)

Gels with the following compositions were prepared by a conventional method.

Ingredients	Formulation amount (% by mass)		
	Example 12	Example 13	Example 14
Compound represented by formula (4) (Production Example 1)	0.1	-	-
Compound represented by formula (5) (Production Example 2)	-	0.1	-
Compound represented by formula (6) (Production Example 3)	-	-	0.1
Decamethylcyclopentasiloxane	10	10	10

Isostearyl isostearate	1	-	-
Olive oil	-	1	-
Macadamia nut oil	-	-	1
Eucalyptus oil	0.1	-	0.1
Hexyldecanol	1	0.1	-
dl- α -tocopherol nicotinate	-	0.1	-
Polyoxyethylene (60) hydrogenated castor oil	2	2	2
Spherical silicone powder *2	1	1	5
Phellodendron bark extract	0.1	1	0.1
Water-soluble chlorophyll	0.02	0.02	0.02
Salvia extract	-	0.3	0.1
1,3-Butylene glycol	5	10	5
Sorbitol liquid	3	3	3
Polyethylene glycol 4000	1	1	1
Carboxyvinyl polymer	0.2	0.2	0.2
Sugar ceramide *3	0.1	0.1	0.1
p-Oxybenzoic acid ester	0.2	0.2	0.2
Mevalonolactone	0.5	0.5	0.5
Edetic acid salt	0.02	0.02	0.02
Potassium hydroxide	0.05	0.05	0.05
Purified water	balance	balance	balance

*2: Tospearl, manufactured by GE Toshiba Silicone Co., Ltd.

*3: Bioceramide, manufactured by Kibun Food Chemical Co., Ltd.

Examples 15 to 17 (Lipophilic cream)

- 5 Lipophilic creams with the following compositions
were prepared by a conventional method.

Ingredients	Formulation amount (% by mass)		
	Example 15	Example 16	Example 17
Compound represented by formula (4) (Production Example 1)	0.1	-	-
Compound represented by formula (5)	-	0.1	-

(Production Example 2)			
Compound represented by formula (6) (Production Example 3)	-	-	0.1
Co-modified silicone *4	2	2	2
Polyeoxethylene-modified silicone dispersion *5	-	2	-
Squalane	-	-	10
Decamethylcyclopentasiloxane	15	20	10
Methylpolysiloxane	5	2	3
Cholesteryl branched long chain fatty acid ester *6	-	-	3
Silicone elastomer dispersion *7	5	2	-
Phellodendron bark extract	1	1	1
Glycyrrhiza extract	0.1	0.1	0.1
Water-soluble chlorophyll	0.02	0.02	0.02
Sodium chloride	1	1	1
Dipropylene glycol	5	5	5
Concentrated glycerin	5	5	5
Raffinose	1	1	1
p-Oxybenzoic acid ester	0.3	0.3	0.3
N-Methyl-L-serine	0.5	0.5	0.5
Purified water	balance	balance	balance

*4: ABIL EM90, manufactured by Goldschmidt A.G.

*5: Silicone BY 22-008, manufactured by Dow Corning Toray Silicone

*6: YOFCO CLE-NH, manufactured by Nippon Fine Chemical Co., Ltd.

*7: Trefil, manufactured by Dow Corning Toray Silicone Co., Ltd.

5

Examples 18 to 20 (Sunscreen)

Sunscreens with the following compositions were prepared by a conventional method.

Ingredients	Formulation amount (% by mass)		
	Example 18	Example 19	Example 20
Compound represented by formula (4) (Production Example 1)	0.1	-	-

Compound represented by formula (5) (Production Example 2)	-	0.1	-
Compound represented by formula (6) (Production Example 3)	-	-	0.1
Diocetyl ether	10	10	10
Co-modified silicone *4	2	2	2
Glyceryl tri-2-ethylhexanoate	5	5	5
Hydrogenated oil	0.1	0.1	0.1
Methylphenylpolysiloxane	3	3	3
Phytosteryl macadamia nut fatty acid	-	-	2
2-Ethylhexyl p-methoxycinnamate	-	7	7
Titanium oxide	5	5	4
Zinc oxide	5	5	4
Phellodendron bark extract	1	1	1
Magnesium chloride	1	1	1
1,3-Butylene glycol	5	5	5
Phenoxyethanol	0.3	0.3	0.3
Hibiscus extract	1	1	1
Aloe extract	0.1	0.1	0.1
Yeast extract *8	1	1	1
Purified water	balance	balance	balance

*4: ABIL EM90, manufactured by Goldschmidt A. G.

*8: Dismutin, manufactured by PentaFarm Ltd.

Example 21 (Face lotion)

Ingredients	Formulation amount (% by mass)
Ethanol	10
Polyoxyethylene (60) hydrogenated castor oil	1
Glycerin	3
1,3-Butylene glycol	2
Dipropylene glycol	3
Polyethylene glycol 1500	1
Phosphoric acid salt	q.s.

Edetic acid salt	q.s.
Methylparaben	q.s.
Compound represented by formula (10) (Production Example 7)	0.1
Anti-oxidant	q.s.
Purified water	balance

Examples 22 and 23 (Emulsion)

Ingredients	Formulation amount (% by mass)	
	Example 22	Example 23
Stearic acid	1	1
Stearic acid glycerin ester	2	2
Cetanol	1	1
Cholesterol	0.5	0.5
Vaseline	2	2
Squalane	5	5
Liquid paraffin	5	5
Silicone oil	1	1
Acylglutamic acid salt	1	1
Xanthane gum	0.5	0.5
Glycerin	2	2
Dipropylene glycol	3	3
Compound represented by formula (5) (Production Example 2)	0.1	-
Compound represented by formula (10) (Production Example 7)	-	0.1
Butylparaben	q.s.	q.s.
Anti-oxidant	q.s.	q.s.
Purified water	balance	balance

Examples 24 and 25 (Cream)

Ingredients	Formulation amount (% by mass)	
	Example 24	Example 25

Stearic acid	2	2
Stearic acid glycerin ester	2	2
Cetanol	3	3
Cholesterol	0.5	0.5
Vaseline	2	2
Squalane	5	5
Liquid paraffin	10	10
Silicone oil	1	1
Acylglutamic acid salt	1	1
Xanthane gum	0.5	0.5
Glycerin	5	5
Dipropylene glycol	3	3
Compound represented by formula (5) (Production Example 2)	0.1	-
Compound represented by formula (10) (Production Example 7)	-	0.1
Butylparaben	q.s.	q.s.
Anti-oxidant	q.s.	q.s.
Purified water	balance	balance

Examples 26 and 27 (Sunscreen)

Ingredients	Formulation amount (% by mass)	
	Example 26	Example 27
Ethanol	10	10
Octyl methoxycinnamate	7	7
POE-POP-modified dimethylpolysiloxane	2	2
Ultrafine titanium oxide particles	5	5
Zinc oxide	5	5
Cyclic silicone	10	10
Diemethylpolysiloxane (6cs)	10	10
Compound represented by formula (5) (Production Example 2)	0.1	-
Compound represented by formula (10) (Production Example 7)	-	0.1

Anti-oxidant	q.s.	q.s.
Purified water	balance	balance

Example 28 (Skin cream)

A skin cream with the following composition was prepared by a conventional method.

Ingredients	Formulation amount (% by mass)
Beeswax	2.0
Stearic acid	5.0
Stearyl alcohol	5.0
Hydrogenated lanolin	2.0
Squalene	20.0
Sorbitan monostearate	3.0
Polyoxyethylene (20) sorbitan monostearate	3.0
Propylene glycol	5.0
Methylparaben	0.2
Compound represented by formula (17)	0.1
Purified water	Balance in the total 100

5

Example 29 (Skin cream)

A skin cream with the following composition was prepared by a conventional method.

Ingredients	Formulation amount (% by mass)
Beeswax	2.0
Stearic acid	5.0
Stearyl alcohol	5.0
Hydrogenated lanolin	2.0
Squalene	20.0
Sorbitan monostearate	3.0
Polyoxyethylene (20) sorbitan monostearate	3.0
Propylene glycol	5.0

Methylparaben	0.2
Compound represented by formula (12) (Production Example 9)	0.5
Purified water	Balance in the total 100

Example 30 (Skin lotion)

A skin lotion with the following composition was prepared by a conventional method.

Ingredients	Formulation amount (% by mass)
Olive oil	10.0
Isopropyl myristate	1.0
Polyoxyethylene (6) nonylphenyl ether	0.5
Propylene glycol	1.0
Glycerin	2.0
Methylparaben	0.1
Ethanol	7.0
Compound represented by formula (17)	0.5
Purified water	Balance in the total 100

5

Example 31 (Bath agent)

A bathing agent with the following composition was prepared by a conventional method.

Ingredients	Formulation amount (% by mass)
Compound represented by formula (17)	1.0
Sodium hydrogen carbonate	balance
Sodium carbonate	20.0
Sodium sulfate	15.0
Sodium chloride	7.5
Silicic acid anhydride	0.5
1,3-butylene glycol	1.0
Urea	1.0
Seaweed extract	1.0
Color additive	q.s.

Dextrin	q.s.
Fragrance	q.s.

Examples 32 to 33 (Skin cream)

Skin creams with the following compositions were prepared by a conventional method.

Ingredients	Formulation amount (% by mass)	
	Example 32	Example 33
Compound represented by formula (17)	0.1	-
Compound represented by formula (14) (Production Example 11)	-	0.1
Stearic acid	1	-
Isostearic acid	-	1
Monostearic acid glycerin	2	2
Behenyl alcohol	2	2
Bleached beeswax	1	1
Cetyl myristate	1	1
Sorbitan sesquioleate	1	1
N-Stearoylphytosphingosine	0.1	0.1
Hydrogenated lecithin	0.1	0.1
Plant squalane	5	5
Octyldodecyl myristate	5	5
Phellodendron bark extract	0.1	1
Pyracantha fortuneana fruit extract	0.1	0.3
Water-soluble glycyrrhiza extract	0.1	0.1
1,3-butylene glycol	5	10
Concentrated glycerin	5	5
p-Oxybenzoic acid ester	0.2	0.2
N-Acetylglucosamine oligomer	0.1	0.1
Ascorbic acid phosphate ester magnesium salt	0.1	0.1
Ascorbic acid phosphate ester sodium salt	0.1	0.1
γ -Aminobutyric acid	0.1	0.1

Sodium N-stearoylglutamate	0.2	0.2
Alkyl-modified carboxyvinyl polymer *1	0.05	0.05
Nicotinic acid amide	0.1	0.1
Sarcosine	0.1	0.1
Purified water	balance	balance

*1: PEMULEN TR-1, manufactured by B. F. Goodrich

Examples 34 and 35 (Lotion)

5 Lotions with the following compositions were
prepared by a conventional method.

Ingredients	Formulation amount (% by mass)	
	Example 34	Example 35
Compound represented by formula (17)	0.1	-
Compound represented by formula (12) (Production Example 9)	-	0.1
Phellodendron bark extract	0.1	0.3
Hibiscus extract	0.2	0.5
Lactic acid bacteria culture medium	0.1	0.1
1,3-Butylene glycol	5	5
Dipropylene glycol	5	5
Raffinose	1	1
Ethanol	1	1
Phenoxyethanol	0.2	0.2
Pectin	0.05	0.05
Xanthane gum	0.1	0.1
Sodium citrate	0.05	0.05
Equisetum arvense extract	0.1	0.1
Diisopropylamine dichloroacetic acid	0.2	0.2
γ -Amino- β -hydroxybutyric acid	0.2	0.2
Sodium hyaluronate	0.001	0.001
Dipotassium glycyrrhizate	0.02	0.02
Naematoloma sublateritium extract	0.05	0.05

Decarboxycarnosine hydrochloride	0.05	0.05
Fragrance	0.02	0.02
Purified water	balance	balance

Examples 36 and 37 (Gel)

Gels with the following compositions were prepared by a conventional method.

Ingredients	Formulation amount (% by mass)	
	Example 36	Example 37
Compound represented by formula (13) (Production Example 10)	0.1	-
Compound represented by formula (15) (Production Example 12)	-	0.1
Decamethylcyclopentasiloxane	10	10
Isostearyl isostearate	1	-
Olive oil	-	1
Macadamia nut oil	0.1	0.1
Eucalyptus oil	0.1	-
Hexyldecanol	1	0.1
dl- α -tocopherol nicotinate	-	0.1
Polyoxyethylene (60) hydrogenated castor oil	2	2
Spherical silicone powder *2	1	1
Phellodendron bark extract	0.1	1
Water-soluble chlorophyll	0.02	0.02
Salvia extract	-	0.3
1,3-Butylene glycol	5	10
Sorbitol liquid	3	3
Polyethylene glycol 4000	1	1
Carboxyvinyl polymer	0.2	0.2
Sugar ceramide *3	0.1	0.1
p-Oxybenzoic acid ester	0.2	0.2
Mevalonolactone	0.5	0.5
Edetic acid salt	0.02	0.02

Potassium hydroxide	0.05	0.05
Purified water	balance	balance

*2: Tospearl, manufactured by GE Toshiba Silicone Co., Ltd.

*3: Bioceramide, manufactured by Kibun Food Chemical Co., Ltd.

Examples 38 and 39 (Lipophilic cream)

- 5 Lipophilic creams with the following compositions were prepared by a conventional method.

Ingredients	Formulation amount (% by mass)	
	Example 38	Example 39
Compound represented by formula (17)	0.1	-
Compound represented by formula (15) (Production Example 12)	-	0.1
Co-modified silicone *4	2	2
Polyeoxethylene-modified silicone dispersion *5	-	2
Squalane	2	2
Decamethylcyclopentasiloxane	15	20
Methylpolysiloxane	5	2
Cholesteryl branched long chain fatty acid ester *6	1	1
Silicone elastomer dispersion *7	5	2
Phellodendron bark extract	1	1
Glycyrrhiza extract	0.1	0.1
Water-soluble chlorophyll	0.02	0.02
Sodium chloride	1	1
Dipropylene glycol	5	5
Concentrated glycerin	5	5
Raffinose	1	1
p-Oxybenzoic acid ester	0.3	0.3
N-Methyl-L-serine	0.5	0.5
Purified water	balance	balance

*4: ABIL EM90, manufactured by Goldschmidt A.G.

*5: Silicone BY 22-008, manufactured by Dow Corning Toray Silicone

*6: YOFCO CLE-NH, manufactured by Nippon Fine Chemical Co., Ltd.

*7: Trefil, manufactured by Dow Corning Toray Silicone Co., Ltd.

Examples 40 and 41 (Sunscreen)

5 Sunscreens with the following compositions were prepared by a conventional method.

Ingredients	Formulation amount (% by mass)	
	Example 40	Example 41
Compound represented by formula (17)	0.1	-
Compound represented by formula (14) (Production Example 11)	-	0.1
Dioctyl ether	10	10
Co-modified silicone *4	2	2
Glyceryl tri-2-ethylhexanoate	5	5
Hydrogenated oil	0.1	0.1
Methylphenylpolysiloxane	3	3
Phytosteryl macadamia nut fatty acid	1	1
2-Ethylhexyl p-methoxycinnamate	-	7
Titanium oxide	5	5
Zinc oxide	5	5
Phellodendron bark extract	1	1
Magnesium chloride	1	1
1,3-Butylene glycol	5	5
Phenoxyethanol	0.3	0.3
Hibiscus extract	1	1
Aloe extract	0.1	0.1
Yeast extract *8	1	1
Purified water	balance	balance

*4: ABIL EM90, manufactured by Goldschmidt A.G.

*8: Dismutin, manufactured by PentaFarm Ltd.

10 Example 42 (Face lotion)

Ingredients	Formulation amount (% by mass)
-------------	-----------------------------------

Ethanol	10
Polyoxyethylene (60) hydrogenated castor oil	1
Glycerin	3
1,3-Butylene glycol	2
Dipropylene glycol	3
Polyethylene glycol 1500	1
Phosphoric acid salt	q.s.
Edetic acid salt	q.s.
Methylparaben	q.s.
Compound represented by formula (15) (Production Example 12)	0.1
Anti-oxidant	q.s.
Purified water	balance

Examples 43 and 44 (Emulsion)

Ingredients	Formulation amount (% by mass)	
	Example 43	Example 44
Stearic acid	1	1
Stearic acid glycerin ester	2	2
Cetanol	1	1
Cholesterol	0.5	0.5
Vaseline	2	2
Squalane	5	5
Liquid paraffin	5	5
Silicone oil	1	1
Acylglutamic acid salt	1	1
Xanthane gum	0.5	0.5
Glycerin	2	2
Dipropylene glycol	3	3
Compound represented by formula (17)	0.1	-
Compound represented by formula (15) (Production Example 12)	-	0.1
Butylparaben	q.s.	q.s.
Anti-oxidant	q.s.	q.s.

Purified water	balance	balance
<hr/>		
Examples 45 and 46 (Cream)		
Ingredients	Formulation amount (% by mass)	
	Example 45	Example 46
Stearic acid	2	2
Stearic acid glycerin ester	2	2
Cetanol	3	3
Cholesterol	0.5	0.5
Vaseline	2	2
Squalane	5	5
Liquid paraffin	10	10
Silicone oil	1	1
Acylglutamic acid salt	1	1
Xanthane gum	0.5	0.5
Glycerin	5	5
Dipropylene glycol	3	3
Compound represented by formula (13) (Production Example 10)	0.1	-
Compound represented by formula (15) (Production Example 12)	-	0.1
Butylparaben	q.s.	q.s.
Anti-oxidant	q.s.	q.s.
Purified water	balance	balance

Examples 47 and 48 (Sunscreen)

Ingredients	Formulation amount (% by mass)	
	Example 47	Example 48
Ethanol	10	10
Octyl methoxycinnamate	7	7
POE-POP-modified dimethylpolysiloxane	2	2
Ultrafine titanium oxide particles	5	5
Zinc oxide	5	5
Cyclic silicone	10	10

Dimethylpolysiloxane (6cs)	10	10
Compound represented by formula (13) (Production Example 10)	0.1	-
Compound represented by formula (15) (Production Example 12)	-	0.1
Anti-oxidant	q.s.	q.s.
Purified water	balance	balance

Industrial Applicability

As described above, it is made clear that the invention can provide an epidermal hyaluronic acid
5 production-promoting agent which can be simply and easily synthesized. The invention makes it possible to prevent the skin from aging (to retain vital, elastic, and moist skin).

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.